ALGEBRAIC GEOMETRY — EXERCISE SHEET 2 DUE ON 10/11/2024

Exercise 0.1. Let $X = \operatorname{Spec} \mathbf{k}[x, y, z]/I \subset \mathbb{A}^3_{\mathbf{k}}$, where $I = (x^2 - yz, y^2 - xz) \subset \mathbf{k}[x, y, z]$. Compute the irreducible components of *X*.

Exercise 0.2. Let *X* be a scheme. Let U, V be affine opens of *X*, and let $x \in U \cap V$. Prove that there exists an affine open neighbourhood *W* of *x* such that *W* is a principal open of both *U* and *V*.

Exercise 0.3. A morphism of schemes $f: X \to Y$ is called *quasicompact* if the preimage of any affine open subset is quasicompact. Prove that $f: X \to Y$ is quasicompact if and only if *Y* has an affine open cover $Y = \bigcup_{i \in I} Y_i$ such that $f^{-1}(Y_i)$ is quasicompact for all *i*.

Exercise 0.4. Assume **k** is an algebraically closed field of characteristic different from 2. Let $Y \subset \mathbb{P}^2_{\mathbf{k}}$ be a nondegenerate conic. Show that *Y* is isomorphic to $\mathbb{P}^1_{\mathbf{k}}$.

Exercise 0.5. Let \mathbb{F} be a field and $f \in \mathbb{F}[x_1, ..., x_n]$ a nonzero polynomial. Show that Spec $\mathbb{F}[x_1, ..., x_n]/(f)$ is reduced (resp. irreducible, resp. integral) if and only if f is square-free (resp. admits only one irreducible factor, resp. is irreducible).

Andrea T. Ricolfi, aricolfi@sissa.it